

Earth gravity field recovery using GPS, GLONASS, and SLR satellites

Krzysztof Sośnica (1), Adrian Jäggi (1),
Daniela Thaller (2), Ulrich Meyer (1),
Gerhard Beutler (1), Rolf Dach (1)

(1) *Astronomical Institute, University of Bern, Switzerland*

(2) *Bundesamt für Kartographie und Geodäsie, Frankfurt am Main, Germany*


18th International Workshop on Laser Ranging

11–15 November 2013 Fujiyoshida, Japan


Motivation

Time-variable Earth's gravity field can be determined from:

- GPS positions of satellites (CHAMP, GRACE-A/B, GOCE, SWARM; high-to-low),
- K-Band GRACE observations (low-to-low),
- Orbit perturbations:
 - using SLR to geodetic satellites,
 - using GPS and GLONASS microwave observations



Sensitivity of GPS resonant orbits



GNSS satellites are very sensitive to gravity field coefficients of degree 2. For coefficients above degree 3, GNSS are only sensitive to **resonant gravity field parameters** (®).

Solution set-up


Estimated parameters		GNSS solutions	SLR solutions
		up to 32 GPS and 24 GLONASS satellites	LAGEOS-1/2, Starlette, Stella, Ajisai
Orbits	Osculating elements	A, e, i, Ω , ω , u_0 (1 set per 3 days)	A, e, i, Ω , ω , u_0 (1 set per 7 days)
	Dynamical parameters	D_0 , Y_0 , X_0 , X_S , X_C (1 set per 3 days)	LAGEOS-1/2: S_0 , S_C , S_S (1 set per 7 days) Sta/Ste/Aji: C_D , S_C , S_S , W_C , W_S (1 set per day)
	Pseudo-stochastic pulses	R, S, W (once per revolution)	LAGEOS-1/2: no pulses Sta/Ste/Aji: S (once per revolution)
Earth rotation parameters		X_P , Y_P , UT1-UTC (1 set per day)	X_P , Y_P , UT1-UTC (1 set per day)
Geocenter coordinates		1 set per 7 days	1 set per 7 days
Earth gravity field		Estimated up to d/o 4/4 (1 set per 7 days)	Estimated up to d/o 4/4 (1 set per 7 days)
Station coordinates		1 set per 7 days	1 set per 7 days
Other parameters		Troposphere ZD (2h), gradients (24h), GNSS-specific translations and ZTD biases	Range biases for selected stations

GNSS solutions are similar to the standard IGS solutions provided by CODE (Center for Orbit Determination in Europe), with some exceptions: Earth gravity field parameters are simultaneously estimated and 7-day instead of 1-day solutions are generated.

SLR solutions are similar to the standard ILRS solutions provided by BKG, but more satellites are included (Sta/Ste/Aji) and Earth gravity field parameters are estimated.

GPS+GLONASS solutions

GNSS orbit modeling





GNSS dynamic orbit parameters estimated in standard CODE solutions:

$$D = D_0$$

$$Y = Y_0$$



$$X = X_0 + X_s \sin \Delta u + X_c \cos \Delta u$$

C_{20} from GPS and GLONASS







GNSS dynamic orbit parameters : D_0, Y_0, X_0, X_s, X_c

The constant and once-per-rev parameters in X are correlated with C_{20}







GNSS dynamic orbit parameters : D_0, Y_0 (no parameters in X)

Zonal spherical harmonics from GPS and GLONASS



Zonal harmonics can be quite well recovered by GNSS




Resonant GPS harmonics



Resonant harmonics, despite a large sensitivity, cannot be fully recovered by GNSS, because of the correlations with D_0 .


SLR solutions

SLR vs. CHAMP vs. GRACE







Some coefficients derived by SLR, CHAMP, and GRACE solutions agree very well. CHAMP solutions show typically larger amplitudes.

SLR vs. CHAMP vs. GRACE







15 out of 21 (71%) coefficients up to d/o 4/4 are derived from SLR with a quality similar to GRACE's
 13 out of 21 (62%) coefficients up to d/o 4/4 are derived from CHAMP with a qual. similar to GRACE's

SLR – specific issues



C_{41} derived by SLR shows similar secular trend to the GRACE results, but the high-frequency part is affected by correlations and modeling deficiencies


SLR – specific issues



Deficiencies in S₂ tide (from the background models) affect not only the GRACE solutions, but also have a minor impact on the SLR solutions.


Low-degree geoid variations [mm]

AIUB-SLR, December 2004




up to d/o 4/4, no filtering

AIUB-GRACE, December 2004




up to d/o 60/45, 1000km Gauss filter

AIUB-GNSS, December 2004



up to d/o 4/4, no filtering

AIUB-CHAMP, December 2004



up to d/o 60/60, 1000km Gauss filter

Low-degree gravity field parameters from SLR solutions fit well to the GRACE results.


Summary

- The gravity field determination using GPS+GLONASS data is very promising, but requires further investigations.
- Most of the low-degree coefficients can be very well established by the observations of SLR geodetic satellites,
- Small issues related to SLR-derived gravity field coefficients originate from:
 - Deficiencies in background models, which are reflected, e.g., in the S_2 alias tide,
 - Deficiencies in the modeling of non-gravitational forces (solar radiation pressure, albedo, the Yarkovsky and Yarkovsky-Schach effects),
 - Correlations between gravity field parameters (e.g., C_{30} and C_{50}) and other parameters (e.g., orbits: perigee, ascending node, etc.).





Etalon-1/2

Thank you for your attention





Geocenter coordinates from GNSS and SLR



Z geocenter component from GNSS is extremely sensitive to orbit modeling; the exclusion of dynamic orbit parameters in the X direction entirely changes the signal!

Geocenter coordinates from GNSS and SLR



References

References:

- Beutler G (2005) Methods of Celestial Mechanics. Volume II: Application to Planetary System, Geodynamics and Satellite Geodesy. Springer Verlag. ISBN 978-3-540-26512-2.
- Dach R, Hugentobler U, Fridez P, Meindl M (2007) Bernese GPS Software Version 5.0. Astronomical Institute, University of Bern, Switzerland.
- Jäggi A, Sośnica K, Thaller D, Beutler G (2012) Validation and estimation of low-degree gravity field coefficients using LAGEOS. Proceedings of 17th ILRS Workshop, Bundesamt für Kartographie und Geodäsie, 48, Frankfurt, 2012
- Michael M, Beutler G, Thaller D, Dach R, Jäggi A (2013) Geocenter coordinates estimated from GNSS data as viewed by perturbation theory. *Advances in Space Research*, Volume 51, Issue 7, p. 1047–1064.
- Rodriguez-Solano CJ, Hugentobler U, Steigenberger P, Lutz S (2012) Impact of Earth radiation pressure on GPS position estimates. *J Geod* 86(5):309–317. doi: 10.1007/s00190-011-0517-4.
- Sośnica K, Thaller D, Dach R, Jäggi A, Beutler G (2013) Impact of atmospheric pressure loading on SLR-derived products and on the consistency between GNSS and SLR results. *Journal of Geodesy* 87 (8), 751–769. DOI: 10.1007/s00190-013-0644-1.
- Sośnica K, Thaller D, Dach R, Jäggi A, Beutler G (2013) Contribution of Starlette, Stella, and AJISAI to the SLR-derived global reference frame. *Journal of Geodesy* (submitted manuscript).
- Sośnica K, Thaller D, Jäggi A, Dach R, Beutler G (2012) Sensitivity of Lageos Orbits to Global Gravity Field Models. *Art Sat*, 47(2), pp. 35–79. doi:10.2478/v10018-012-0013-y.
- Sośnica K, Thaller D, Jäggi A, Dach R, Beutler G (2012) Can we improve LAGEOS solutions by combining with LEO satellites? *Proceedings of the International Technical Laser Workshop 2012 (ITLW-12)*, Frascati (Rome), Italy, November 5–9, 2012.
- Sośnica K, Thaller D, Dach R, Jäggi A, Beutler G (2013) Time variable Earth's gravity field from SLR data. *Advances in Space Research* (to be submitted).
- Sośnica K, Jäggi A, Thaller D, Beutler G, Dach R, Baumann C (2013) SLR-derived terrestrial reference frame using observations to LAGEOS-1/2, Starlette, Stella, and AJISAI. *18th International Workshop on Laser Ranging*, 11–15 November 2013 Fujiyoshida, Japan.
- Thaller D, Sośnica K, Dach R, Jäggi A, Beutler G (2011) LAGEOS-ETALON solutions using the Bernese Software. *Mitteilungen des Bundesamtes fuer Kartographie und Geodäsie*, Proceedings of the 17th International Workshop on Laser Ranging, Extending the Range, Bad Kötzting, Germany, May 16– 20, 2011, vol. 48, pp.333–336, Frankfurt.
- Thaller D, Sośnica K, Mareyen M, Dach R, Jäggi A, Beutler G (2013) Geodetic parameters estimated from LAGEOS and Etalon data and comparison to GNSS–estimates. *Journal of Geodesy* (submitted manuscript).
- Thaller D, Sośnica K, Dach R, Jäggi A, Beutler G, Mareyen M, Richter B (2013) Geocenter coordinates from GNSS and combined GNSS–SLR solutions using satellite co-locations. *IAG Symposia Series* 139.