

# How well can the combination of hISST and SLR replace GRACE?

A discussion from the point of view of applications

M. Weigelt, T. van Dam,  
O. Baur, M.J. Tourian,  
H. Steffen, K. Sośnica,  
A. Jäggi, N. Zehentner,  
T. Mayer-Gürr, N. Sneeuw

RUES

RESEARCH UNIT  
IN ENGINEERING  
SCIENCES

30.09.2014

FACULTY OF SCIENCES, TECHNOLOGY AND COMMUNICATION



LANTMÄTERIET



UNIVERSITÉ DU  
LUXEMBOURG

**AIUB**



# PROCESSING

A short overview

# HLSST + SLR

## High-low SST:

- 11 satellites with GPS

CHAMP	GRACE A	GRACE B
GOCE	MetOp A	MetOp B
TerraSAR X	TanDEM-X	
Swarm A	Swarm B	Swarm C

- accelerometer data used when available
- combination at the normal equation level

# HLSST + SLR

## High-low SST:

- 11 satellites with GPS

CHAMP	GRACE A	GRACE B
GOCE	MetOp A	MetOp B
TerraSAR X	TanDEM-X	
Swarm A	Swarm B	Swarm C

- accelerometer data used when available
- combination at the normal equation level

Details in Zehentner et al:

*Non-dedicated satellite missions for time-variable gravity field recovery*

Today: 14:30

# HLSST + SLR

## High-low SST:

- 11 satellites with GPS

CHAMP	GRACE A	GRACE B
GOCE	MetOp A	MetOp B
TerraSAR X	TanDEM-X	
Swarm A	Swarm B	Swarm C

- accelerometer data used when available
- combination at the normal equation level

## SLR:

- 9 satellites

Lageos 1	Lageos 2	LARES
Starlette	Stella	Larets
AJISAI	Beacon-C	Blits

- gravity parameters together with station coordinates, ERP, geocenter and range biases
- combination at the normal equation level

Details in Zehentner et al:

*Non-dedicated satellite missions for time-variable gravity field recovery*

Today: 14:30

# HLSST + SLR

## High-low SST:

- 11 satellites with GPS

CHAMP	GRACE A	GRACE B
GOCE	MetOp A	MetOp B
TerraSAR X	TanDEM-X	
Swarm A	Swarm B	Swarm C

- accelerometer data used when available
- combination at the normal equation level

Details in Zehentner et al:

*Non-dedicated satellite missions for time-variable gravity field recovery*

Today: 14:30

## SLR:

- 9 satellites

Lageos 1	Lageos 2	LARES
Starlette	Stella	Larets
AJISAI	Beacon-C	Blits

- gravity parameters together with station coordinates, ERP, geocenter and range biases
- combination at the normal equation level

Details in Sośnica et al:

*Time varying gravity from SLR, combined SLR and high-low satellite-to-satellite tracking data*


Today: 14:15

# Data availability




■ dedicated gravity field missions   
 ■ non-dedicated mission   
 ■ spherical SLR   
 ■ non-spherical SLR


# CHAMP-only




# Combined hISST



# Combined hISST vs SLR




# hISST + SLR




# Post-processing: old approach

- Kalman filtering
  - ... allows to handle small numbers of sample (filter warm-up issue).



# Post-processing: old approach


- Kalman filtering
  - ... allows to handle small numbers of sample (filter warm-up issue).




... is equivalent to an AR(2) process: filter design is limited.

# Post-processing: new approach

- Bandpass filtering  
... is possible due to the longer time series (11 years)



# hISST + SLR + post-processing




# MASS TRENDS AND ANNUAL SIGNALS


# Trends

- Total change in terms of EWH
- Gaussian filtering with 750 km


hISST-only



GRACE CSR




hISST & SLR




# Trends

- Total change in terms of EWH
- Gaussian filtering with 750 km


hISST-only



GRACE CSR



hISST & SLR



# Trends

Mass change rates in [Gt/yr] of areas with the strongest signals

Region	hISST-only <sup>a</sup>		hISST + SLR		GRACE CSR <sup>a</sup>
		$\Delta$ Grace		$\Delta$ Grace	
Greenland	$-252 \pm 10$	-12%	$-267 \pm 12$	-6%	$-285 \pm 10$
Greenland ext. <sup>b</sup>	$-275 \pm 12$	-13%	$-294 \pm 12$	-7%	$-316 \pm 10$
Canadian Shield	$152 \pm 10$	-12%	$158 \pm 6$	-8%	$172 \pm 6$
West Antarctica	$-127 \pm 10$	-10%	$-119 \pm 10$	-15%	$-140 \pm 10$
East Antarctica	$103 \pm 10$	0%	$104 \pm 10$	1%	$104 \pm 6$

<sup>a</sup>  $C_{20}$  replaced by values from SLR.

<sup>b</sup> including Iceland, Svalbard, and the Canadian Arctic archipelago

Uncertainties are give at the 95% ( $2\sigma$ ) confidence level.

# Trends

Mass change rates in [Gt/yr] of areas with the strongest signals

Region	hISST-only <sup>a</sup>		hISST + SLR		GRACE CSR <sup>a</sup>
		$\Delta$ Grace		$\Delta$ Grace	
Greenland	$-252 \pm 10$	-12%	$-267 \pm 12$	-6%	$-285 \pm 10$
Greenland ext. <sup>b</sup>	$-275 \pm 12$	-13%	$-294 \pm 12$	-7%	$-316 \pm 10$
Canadian Shield	$152 \pm 10$	-12%	$158 \pm 6$	-8%	$172 \pm 6$
West Antarctica	$-127 \pm 10$	-10%	$-119 \pm 10$	-15%	$-140 \pm 10$
East Antarctica	$103 \pm 10$	0%	$104 \pm 10$	1%	$104 \pm 6$

<sup>a</sup>  $C_{20}$  replaced by values from SLR.


<sup>b</sup> including Iceland, Svalbard, and the Canadian Arctic archipelago

Uncertainties are give at the 95% ( $2\sigma$ ) confidence level.


# Annual signal

Gaussian filtering with 750 km

## CHAMP-only




## GRACE CSR




# Annual signal

Gaussian filtering with 750 km


## CHAMP-only



## hISST-only




## GRACE CSR




# Annual signal

Gaussian filtering with 750 km


CHAMP-only




hISST-only






GRACE CSR



hISST & SLR



# Annual signal




# HYDROLOGY AND HYDRO-METEOROLOGY

# Mass change as a hydrological observable

$ET_a$  = Evapotranspiration (MODIS)

$P$  = Precipitation (GPCC)

$R$  = Runoff (GRDC)



$\nabla \cdot \bar{Q}$  = divergence of vertically integrated moisture flux (ERA-interim)




# Hydrological validation: $P-ET_a-R$

Gaussian  
filtering with  
750km

Correlation:

GRACE




hISST & SLR



RMS ratio:

GRACE

hISST & SLR






# Hydro-meteorological validation: $\nabla \cdot Q - R$

Gaussian filtering with 750km

Correlation:

GRACE




hISST & SLR




RMS ratio:

GRACE

hISST & SLR




# Hydrology and hydro-meteorology



# GLACIAL ISOSTATIC ADJUSTMENT

# North America

## Gaussian filtering with 750km



SE maximum:

	$\varphi$	$\lambda$
hISST	49°N	80°W
hISST & SLR	52°N	80°W
CSR	54°N	75°W

W maximum:


	$\varphi$	$\lambda$
hISST	60°N	99°W
hISST & SLR	58°N	100°W
CSR	63°N	99°W

# Fennoscandia

Gaussian filtering with 750km

Maximum:

	$\varphi$	$\lambda$
hiSST	65°N	18°W
hiSST & SLR	65°N	18°W
CSR	63°N	20°W



# HOW GOOD ARE WE THEN?

# What can be done:

# What can be done:

- Spatial resolution for the temporally filtered and spatially unfiltered signal is between 1000 to 2000 km
  - ... can be improved by spatial filtering – e.g. Gaussian 750km

# What can be done:

- Spatial resolution for the temporally filtered and spatially unfiltered signal is between 1000 to 2000 km
  - ... can be improved by spatial filtering – e.g. Gaussian 750km
- Mass trend and annual signals are approximately 5-15% underestimated.

# What can be done:

- Spatial resolution for the temporally filtered and spatially unfiltered signal is between 1000 to 2000 km
  - ... can be improved by spatial filtering – e.g. Gaussian 750km
- Mass trend and annual signals are approximately 5-15% underestimated.
- Hydrological signal is similarly underestimated in wet areas. In arid areas the signal is overestimated (SNR!)

# What can be done:

- Spatial resolution for the temporally filtered and spatially unfiltered signal is between 1000 to 2000 km
  - ... can be improved by spatial filtering – e.g. Gaussian 750km
- Mass trend and annual signals are approximately 5-15% underestimated.
- Hydrological signal is similarly underestimated in wet areas. In arid areas the signal is overestimated (SNR!)
- HISST-only and hISST+SLR are able to resolve the double peak of the GIA signal in North America. The smaller Fennoscandia is still overestimated.

# WHAT ARE THE LIMITATIONS?

# Limitations:

# Limitations:

- The methodology is a post-processing method.

# Limitations:

- The methodology is a post-processing method.
- It benefits from the estimation of the mean signals → the longer the time series the better.

# Limitations:

- The methodology is a post-processing method.
- It benefits from the estimation of the mean signals → the longer the time series the better.
- Filtering process smears out episodic events.

# Limitations:

- The methodology is a post-processing method.
- It benefits from the estimation of the mean signals → the longer the time series the better.
- Filtering process smears out episodic events.
- Approach depends 1:1 on the precision of GNSS and SLR observations → spatial resolution remains limited


# Limitations:

- The methodology is a post-processing method.
- It benefits from the estimation of the mean


... but the time series can be continued as satellite data will be available throughout the time to GRACE Follow-On

and SLR observations → spatial resolution remains limited

# GRACE data availability



# hISST + SLR data availability



UNIVERSITÉ DU LUXEMBOURG

# THANK YOU

Home Methodology Downloads Publications

Search:

Home

Methodology

Downloads

Publications

Home

## Welcome to the GEOSAT.UNI.LU

Here we present you results from the gravity field recovery efforts within the [Geophysics Laboratory \(LGP\)](#) at the [University of Luxembourg](#). Please use the menu to the left to go to your area of interest. We present details on the methodology as well as data that is available for download.

# http://geosat.uni.lu

Generally, data that has been published is freely available. Access to other data may be limited and at times special conditions apply. In order to access these data, please send an email to [matthias.weigelt@uni.lu](mailto:matthias.weigelt@uni.lu) in order to get access to the protected area. The [login](#) to the protected area is on the top right.

### About us:

The LGP focuses on four main topics: climate variability, geodynamics, resource management and regional processes. The main objectives are the development of reliable measurement of environmental changes at all spatial and temporal scales, the assessment of the relative impact of human and natural factors in these changes, a better understanding of the relation of the various environmental changes and the evaluation of the risk they induce for mankind. The research relies mainly on the modeling and the interpretation of satellite data.

The LGP is located within the [Research Unit in Engineering Science \(RUES\)](#) at the [Faculté des Sciences, de la Technologie et de la Communication](#), University of Luxembourg. The LGP also hosts the [Global Geophysical Fluid Center \(GGFC\)](#).

UNIVERSITÉ DU LUXEMBOURG

RUES | RESEARCH UNIT IN ENGINEERING SCIENCES

Geophysics

Global Geophysical Fluid Center

University [Legal notice](#)